



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2021

---

## **Coronary CT angiography as a guide in decision-making for left main spontaneous coronary dissection**

Benetos, Georgios ; Afendoulis, Dimitrios ; Buechel, Ronny R ; Smyrnioudis, Nikolaos ; Protogeros, Dimitrios ; Kartalis, Athanasios

DOI: <https://doi.org/10.1016/j.hjc.2020.09.007>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-196074>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Benetos, Georgios; Afendoulis, Dimitrios; Buechel, Ronny R; Smyrnioudis, Nikolaos; Protogeros, Dimitrios; Kartalis, Athanasios (2021). Coronary CT angiography as a guide in decision-making for left main spontaneous coronary dissection. *Hellenic Journal of Cardiology*, 62(4):320-321.

DOI: <https://doi.org/10.1016/j.hjc.2020.09.007>



Contents lists available at ScienceDirect

Hellenic Journal of Cardiology

journal homepage: <http://www.journals.elsevier.com/hellenic-journal-of-cardiology/>

## Correspondence

## Coronary CT angiography as a guide in decision-making for left main spontaneous coronary dissection

## Keywords:


spontaneous coronary dissection  
left main  
CCTA  
CABG

A 36-year-old woman presented to the emergency department with sudden-onset chest pain. Her medical history included an atrial septal defect closure with percutaneous device due to relapsing transient ischemic attacks (TIAs) 7 years prior. A few minutes after presentation, she suffered in-hospital cardiac arrest with ven-

anterior leads. The patient underwent emergent invasive coronary angiography under dual antiplatelet therapy and heparin, which revealed spontaneous coronary dissection of the left main artery extending to the middle segment of the left anterior descending artery (LAD) (Fig. 1, panel A).

The patient was treated conservatively, and her hospitalization was uneventful. She was discharged under treatment with aspirin, clopidogrel, and beta-blocker. She was scheduled for follow-up coronary CT angiography (CCTA) one month after the event, as intubation of coronary ostium and contrast injection during subsequent invasive coronary angiography could potentially lead to further expansion of the dissection.

At presentation one month later, the patient was free of symptoms. CCTA revealed an extensive hematoma at the left main stem, which significantly compromised the true lumen. The dissec-



**Fig. 1.** Panel A: Invasive coronary angiography, revealing left main dissection, extending to the middle LAD; Panel B: First CCTA multiplanar reconstruction (MPR) image of LM and bifurcation, showing the intramural hematoma and the dissection flap; Panel C: Curved MPR image of LAD clearly showing the dissection flap.

tricular fibrillation as underlying rhythm. After successful defibrillation, the electrocardiogram showed ST segment elevation in


tion flap was visible at the level of left main bifurcation (Fig. 1, panel B and C). There was no argument for dissection extending to left anterior descending artery. All three coronary arteries showed no other signs of stenosis.

Peer review under responsibility of Hellenic Society of Cardiology.

<https://doi.org/10.1016/j.hjc.2020.09.007>

1109-9666/© 2020 Hellenic Society of Cardiology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article as: G. Benetos, D. Afendoulis, R.R. Buechel *et al.*, Coronary CT angiography as a guide in decision-making for left main spontaneous coronary dissection, Hellenic Journal of Cardiology, <https://doi.org/10.1016/j.hjc.2020.09.007>



**Fig. 2.** **Panel A:** Second CCTA short axis MPR image at the bifurcation level, showing the dissection flap; **Panel B:** Second CCTA long axis MPR image showing the persisting hematoma and the dissection flap; **Panel C:** Intraoperative image showing intraluminal LAD thrombus proximal to the arteriotomy.

As the patient remained asymptomatic, it was decided after careful consideration to have her undergo a second CCTA two months after the index event. Unfortunately, the second CCTA revealed similar results, with significantly compromised left main lumen due to persisting hematoma and a dissection flap at the level of bifurcation (Fig. 2, panel A and B).

Lastly, taking into consideration the data of the two CCTAs, the Heart Team recommended a surgical revascularization. Indeed, the patient underwent coronary artery bypass surgery two and a half months after the index event. Both left and right internal mammary arteries (LIMA and RIMA) were harvested. LAD arteriotomy was performed before the bifurcation to the first diagonal branch. The dissection with superimposing thrombus (2 cm, Fig. 2, panel C) was visible proximally, and the thrombus was removed appropriately. The true LAD lumen was identified, and RIMA-LAD anastomosis was completed successfully with a prolene 7-0 suture. Obtuse marginalis 1 (OM1) showed no signs of dissection, and LIMA was successfully anastomosed with a prolene-7 suture. The post-operative course was uneventful, and the patient was discharged on the sixth post-operative day.

CCTA is a valuable tool for noninvasive angiographic follow-up of patients with spontaneous dissection.<sup>1-3</sup> Although conservative management is mostly proposed, data on appropriate management of stabilized and further asymptomatic patients with left main dissection remain extremely scarce. In the present case, serial imaging with CCTA contributed significantly in decision making.

## Disclosures

There is no conflict of interest to disclose for any author.

## References

1. Griffin A, Navar AM, Crawford L, Mammarappallil J, Koweek LH. A case of spontaneous coronary artery dissection diagnosed by coronary computed tomography angiography. *J Cardiovasc Comput Tomogr.* 2018;12:88–89.
2. Roura G, Ariza-Sole A, Rodriguez-Caballero IF, et al. Noninvasive Follow-Up of Patients With Spontaneous Coronary Artery Dissection With CT Angiography. *JACC Cardiovasc Imaging.* 2016;9:896–897.
3. Gad MM, Mahmoud AN, Saad AM, et al. Incidence, Clinical Presentation, and Causes of 30-Day Readmission Following Hospitalization With Spontaneous Coronary Artery Dissection. *JACC Cardiovasc Interv.* 2020;13:921–932.

Georgios Benetos\*

First Department of Cardiology, Athens School of Medicine, Greece

Dimitrios Afendoulis

Cardiology Clinic, General Hospital of Chios, Greece

Ronny R. Buechel

Department of Nuclear Medicine, University Hospital of Zurich, Switzerland

Nikolaos Smyrnioudis

Cardiology Clinic, General Hospital of Chios, Greece

Dimitrios Protogeros

Department of Cardiothoracic Surgery, Henry Dynant Hospital, Athens, Greece

Athanasios Kartalis

Cardiology Clinic, General Hospital of Chios, Greece

\* Corresponding author: Georgios Benetos, MD, First Department of Cardiology, Athens School of Medicine, 194, Aleksandras Av., 11521, Athens, Greece. Phone: +302106426812. E-mail address: [benetosg@gmail.com](mailto:benetosg@gmail.com) (G. Benetos).

20 August 2020

Available online xxx